

REMARKS**I. Status of the Claims**

Claims 1-8 are withdrawn.

Claims 10, 11, 13 and 14 are amended.

Claims 16 and 17 are new.

Claims 9-17 are pending.

II. Interview Summary

Applicant thanks Supervisor Padmanabhan and Examiner Cotton for suggestions on claim amendments to move this case toward allowance. A question of total daily dosage as distinct from unit dose, e.g. dose per capsule, arose. Claim amendments with support from the specification were suggested, as long as the claim scope was not within the art, e.g. in Examples 1-46 in Caruso. Support for the daily doses and unit doses are found on at least the following locations. Citations are to paragraphs in the specification:

What is a "low dose" tricyclic antidepressant?

[00008] 25 mg/day or less.

[00010] 0.5 gm-2.6 gm daily; 0.5-2 gm/day acetaminophen; 0.6-2.6 gm/day aspirin; 0.6-1.8 gm/day ibuprofen.

What is the "standard dose" of non-narcotic analgesic?

[00011] 2.5 mg to 25 mg/day (.5 mg to 2 mg, 10-15 mg/day).

[00015] Example 1: 500 mg acetaminophen + 5 mg doxepin (unit dose).

[00016] Example 2: 5 mg doxepin + 650 mg aspirin, unit dose, twice daily.

[00017] Example 3: 10 mg doxepin + 600 mg ibuprofen.

The references requested by the examiner regarding low and standard doses are in Exhibits A and B.

No fees are believed due at this time, however, please charge any additional deficiencies or credit any overpayments to deposit account number 12-0913 with reference to our attorney docket number (41957/102748).

Respectfully submitted,

A handwritten signature in cursive script, appearing to read "Alice O. Martin".

Alice O. Martin
Registration No. 35,601


February 12, 2007
Barnes & Thornburg LLP
P.O. Box 2786
Chicago, IL 60690-2786

14TH
EDITION

Harrison's

**PRINCIPLES of
INTERNAL
MEDICINE**

**Fauci
Braunwald
Isselbacher
Wilson
Martin
Kasper
Hauser
Longo**



BEST AVAILABLE COPY



Note: Dr. Fauci and Dr. Longo's works as editors and authors were performed outside the scope of their employment as U.S. government employees. These works represent their personal and professional views and not necessarily those of the U.S. government.

**Harrison's
PRINCIPLES OF INTERNAL MEDICINE
Fourteenth Edition**

Copyright © 1998, 1994, 1991, 1987, 1983, 1980, 1977, 1974, 1970, 1966, 1962, 1958 by *The McGraw-Hill Companies, Inc.* All rights reserved. Printed in the United States of America. Except as permitted under the United States Copyright Act of 1976, no part of this publication may be reproduced or distributed in any form or by any means, or stored in a data base or retrieval system, without the prior written permission of the publisher.

4567890 DOWDOW 0403020100
ISBN 0-07-020291-5 (COMBO)
0-07-020292-3 (VOL 1)
0-07-020293-1 (VOL 2)
0-07-912013-X (SET)

FOREIGN LANGUAGE EDITIONS

Arabic (Thirteenth Edition)—McGraw-Hill Libri Italia srl (est. 1996)
Chinese (Twelfth Edition)—McGraw-Hill Book Company—Singapore © 1994
Croatian (Thirteenth Edition)—McGraw-Hill Libri Italia srl (est. 1996)
French (Thirteenth Edition)—McGraw-Hill Libri Italia srl © 1995
German (Thirteenth Edition)—McGraw-Hill Libri Italia srl © 1995
Greek (Twelfth Edition)—Parissianos © 1994
Italian (Thirteenth Edition)—McGraw-Hill Libri Italia srl © 1995
Japanese (Eleventh Edition)—Hirokawa © 1991
Portuguese (Thirteenth Edition)—McGraw-Hill/Nueva Editorial Interamericana © 1995
Spanish (Thirteenth Edition)—McGraw-Hill Interamericana de España © 1994
Turkish (Thirteenth Edition)—McGraw-Hill Libri Italia srl (est. 1996)

This book was set in Times Roman by Monotype Composition Company. The editors were J. Dereck Jeffers, Martin J. Wonsiewicz, and Mariapaz Ramos Englis; editorial assistants were Rose Derario and Daniel J. Green. The production director was Robert Laffler. The index was prepared by Irving Tullar. The text and cover designer was Marsha Cohen/Parallelogram Graphics.

R.R. Donnelley and Sons, Inc., was the printer and binder.

Library of Congress Cataloging-in-Publication Data

Harrison's principles of internal medicine.—14th. ed./editors,
Anthony S. Fauci . . . [et al.]
p. cm.
Includes bibliographical references and index.
ISBN 0-07-020291-5 (1 vol. ed.).—ISBN 0-07-912013-X (2 vol. ed.
: set).—ISBN 0-07-020292-3 (2 vol. ed. : bk 1).—ISBN
0-07-020293-1 (2 vol. ed. : bk 2)
1. Internal medicine. I. Harrison, Tinsley Randolph.
II. Fauci, Anthony S.
[DNLM: 1. Internal Medicine. WB 115 H322 1998]
RC46.H333 1998
616—dc21
DNLM/DLC
for Library of Congress

97-14184

BEST AVAILABLE COPY

The introduction of a parenteral form of NSAID, ketorolac, extends the usefulness of this class of compounds in the management of acute severe pain. Ketorolac is sufficiently potent and rapid in onset to supplant opioids for many patients with acute severe headache and musculoskeletal pain.

Opioid Analgesics Opioids are the most potent pain-relieving drugs currently available. Furthermore, of all analgesics, they have the broadest range of efficacy, providing the most reliable method for rapidly relieving pain. Although side effects are common, except for respiratory depression, they are usually not serious and can be reversed rapidly with the narcotic antagonist naloxone. The physician should not hesitate to use opioid analgesics in patients with acute severe pain. Table 12-1 lists the most commonly used opioid analgesics.

Opioids produce analgesia by actions in the central nervous system. They activate pain-inhibitory neurons and directly inhibit pain-transmission neurons. Most of the commercially available opioid analgesics act at the same opiate receptor (mu receptor), differing mainly in potency, speed of onset, duration of action, and optimal route of administration. Although the dose-related side effects (sedation, respiratory depression, pruritus, constipation) are similar among the different opioids, some side effects are due to accumulation of nonopioid metabolites that are unique to individual drugs. One striking example of this is normeperidine, a metabolite of meperidine. Normeperidine produces hyperexcitability and seizures that are not

reversible with naloxone. Normeperidine accumulation is greater in patients with renal failure.

The most rapid relief with opioids is obtained by intravenous administration; relief with oral administration is significantly less. Common acute side effects include nausea, vomiting, and sedation. These effects are dose-related, and there is great variability among patients in the doses that relieve pain and produce side effects. Because of this, initiation of therapy requires titration to optimal dose and interval. The most important principle is to provide adequate pain relief. This requires asking the patient whether the drug has relieved the pain and, if so, when the relief wears off. The most common error made by physicians in managing severe pain with opioids is to prescribe an inadequate dose. Since many patients are reluctant to complain, this practice leads to needless suffering. In the absence of sedation at the expected time of peak effect, a physician should not hesitate to repeat the initial dose to achieve satisfactory pain relief.

An innovative approach to the problem of achieving adequate pain relief is the use of patient-controlled analgesia (PCA). PCA requires a device that immediately delivers a pre-programmed dose of an opioid drug when the patient pushes a button. The device can be programmed to limit the total hourly dose so that overdosing is impossible. The patient can then titrate the dose to the optimal level. This approach is used most extensively for the management of postoperative pain, but there is no reason why it should not be used for any hospitalized patient with persistent severe pain. PCA can also be used for home care of patients with intractable pain, such as metastatic cancer.

Table 12-1

Drugs for Relief of Pain

NONNARCOTIC ANALGESICS: USUAL DOSES AND INTERVALS

Generic Name	Dose, mg	Interval	Comments
Acetylsalicylic acid	650 PO	q 4 h	Enteric-coated preparations available
Acetaminophen	650 PO	q 4 h	Side effects uncommon
Ibuprofen	400 PO	q 4-6 h	Available without prescription
Naproxen	250-500 PO	q 12 h	Delayed effects may be due to long half-life
Fenoprofen	200 PO	q 4-6 h	pa
Indomethacin	25-50 PO	q 8 h	Gastrointestinal side effects common
Ketorolac	15-60 IM	q 4-6 h	Available for parenteral use (IM)

NARCOTIC ANALGESICS: USUAL DOSES AND INTERVALS

Generic Name	Parenteral Dose, mg	PO Dose, mg	Comments
Codeine	30-60 q 4 h	30-60 q 4 h	Nausea common
Oxycodone	—	5-10 q 4-6 h	Usually available with acetaminophen or aspirin
Morphine	10 q 4 h	60 q 4 h	
Morphine sustained release	pa	60-180 bid to tid	Oral slow-release preparation
Hydromorphone	1-2 q 4 h	2-4 q 4 h	Shorter acting than morphine sulfate
Levorphanol	2 q 6-8 h	4 q 6-8 h	Longer acting than morphine sulfate; absorbed well PO
Methadone	10 q 6-8 h	20 q 6-8 h	Delayed sedation due to long half-life
Meperidine	75-100 q 3-4 h	300 q 4 h	Poorly absorbed PO; normeperidine a toxic metabolite
Butorphanol	—	1-2 q 4 h	Intranasal spray
Fentanyl	—	—	Transdermal patch

ANTICONVULSANTS AND ANTIARRHYTHMICS

Generic Name	PO Dose, mg	Interval
Phenytoin	300	daily/qhs
Carbamazepine	200-300	q 6 h
Clonazepam	1	q 6 h
Mexiletine	150-300	q 6-12 h

TRICYCLIC ANTIDEPRESSANTS

Generic Name	Uptake Blockade		Sedative Potency	Anticholinergic Potency	Orthostatic Hypotension	Cardiac Arrhythmia	Average Dose, mg/day	Range, mg/day
	SHT	NE						
Doxepin	++	+	High	Moderate	Moderate	Less	200	75-400
Amiripryline	++++	++	High	Highest	Moderate	Yes	150	25-300
Imipramine	++++	++	Moderate	Moderate	High	Yes	200	75-400
Nortriptyline	+++	++	Moderate	Moderate	Low	Yes	100	40-150
Desipramine	+++	++++	Low	Low	Low	Yes	150	50-300

The
PHARMACOLOGICAL
BASIS OF
THERAPEUTICS

Joel G. Hardman
Lee E. Limbird

Perry B. Molinoff
Raymond W. Ruddon

Alfred Goodman Gilman

BEST AVAILABLE COPY

McGraw-Hill

A Division of The McGraw-Hill Companies



Goodman and Gilman's THE PHARMACOLOGICAL BASIS OF THERAPEUTICS, 9/e

Copyright © 1996, 1990, 1985, 1980, 1975, 1970, 1965, 1955, 1941 by The McGraw-Hill Companies, Inc. All rights reserved. Printed in the United States of America. Except as permitted under the United States Copyright Act of 1976, no part of this publication may be reproduced or distributed in any form or by any means, or stored in a data base or retrieval system, without the prior written permission of the publisher.

234567890 DOWDOW 9987

ISBN 0-07-026266-7

This book was set in Times Roman by York Graphic Services, Inc. The editors were Martin J. Wonsiewicz and Peter McCurdy; the production supervisors were Robert Laffler and Clare Stanley, and the cover designer was Marsha Cohen/Paralellogram. The index was prepared by Irving Condé Tullar.

R.R. Donnelley and Sons Company was printer and binder.

This book is printed on acid-free paper.

Library of Congress Cataloging-in-Publication Data

Goodman & Gilman's *The Pharmacological Basis of Therapeutics*. —9th ed. / Joel G. Hardman, Alfred Goodman Gilman, Lee E. Limbird.

p. cm.

Includes bibliographical references and index.

ISBN 0-07-026266-7 (hardcover)

I. Pharmacology. 2. Chemotherapy. I. Goodman, Louis Sanford. II. Gilman, Alfred. III. Hardman, Joel G. IV. Gilman, Alfred Goodman. V. Limbird, Lee E.

[DNLM: 1. Pharmacology. 2. Drug Therapy. QV 4 G6532 1995]

RM300.G644 1995

615'.7—dc20

DNLM/DLC

for Library of Congress

95-36658

BEST AVAILABLE COPY

ler in the
enzyl de-
sics, and
dibenz-
only by
a seven-
psychotic
few com-
sedative






analogs,
hiazines,
psychotic
patients;
has accu-

l its iso-
ment of
ig effects
rs found
siting the
gues and
patients
t of mod-

Tricyclic
ically to
on clini-
es, imip-
tabolite)
ne, there
riptyline
oxepine)
itally re-
ne (con-
x-carbon
tidepres-
a three-
in most
lic anti-
e agents
of nor-

and other
h diben-
ally, the
lar stere-
rings, as
gener of
tidepres-
1 below.
agent re-
more ef-
fizations
and nor-
thioxan-
e chem-
ents are

Table 19-1
Antidepressants: Chemical Structures, Dose and Dosage Forms, and Side Effects

NONPROPRIETARY NAME (TRADE NAME)			DOSE AND DOSAGE FORMS			SIDE EFFECTS						
Norepinephrine-Reuptake Inhibitors: Tertiary Amine Tricyclics 			<i>Usual Dose, mg/day</i>	<i>Extreme Dose, mg/day</i>	<i>Dosage Form</i>	<i>Amine Effects</i>	<i>Sedation</i>	<i>Anti-cholinergic Effects</i>	<i>Hypo-tension</i>	<i>Cardiac Effects</i>	<i>Seizures</i>	<i>Weight Gain</i>
<i>R</i> ₁	<i>R</i> ₂	<i>R</i> ₃										
Amiripryline (ELAVIL and others)												
C	H	C=CH(CH ₂) ₂ N(CH ₃) ₂	100-200	25-300	O, I	NE, 5-HT	+++	+++	+++	+++	++	++
Clomipramine (ANAFRANIL)												
C	Cl	N-(CH ₂) ₃ N(CH ₃) ₂	100-200	25-250	O	NE, 5-HT	++	+++	++	+++	+++	+
Doxepin (ADAPIN, SINEQUAN)												
O	H	N=CH(CH ₂) ₂ N(CH ₃) ₂	100-200	25-300	O	NE, 5-HT	+++	++	+++	++	++	++
Imipramine (TOFRANIL and others)												
C	H	N-(CH ₂) ₃ N(CH ₃) ₂	100-200	25-300	O, I	NE, 5-HT	++	++	++	+++	++	++
(+)-Trimipramine (SURMONTIL)												
												
C	H	N-CH ₂ CH(CH ₃) ₂	75-200	25-300	O	NE, 5-HT	+++	+++	++	+++	++	++
Norepinephrine-Reuptake Inhibitors: Secondary Amine Tricyclics Anoxapine (ASENDIN)												
			200-300	50-600	O	NE, DA	+	+	++	++	++	+
Desipramine (NORPRAMIN, PERTOFRANE)												
			100-200	25-300	O	NE	0/+	+	+	++	+	+
Maprotiline (LUDIOMIL)												
			100-150	25-225	O	NE	++	++	++	++	+++	+